

J6673(C)  
Y2-R566-EDG

**SKIN CARE PRODUCT CONTAINING RETINOIDS, RETINOID BOOSTER AND  
PHYTOESTROGENS IN A DUAL COMPARTMENT PACKAGE**

This application claims priority under 35 U.S.C. § 119 from provisional  
5 application Serial No. 60/258,457, filed December 28, 2000.

**FIELD OF THE INVENTION**

The invention relates to stable skin care compositions containing a retinoid in  
10 a first compartment and a retinoid booster system and a phytoestrogen in a second  
compartment of a dual compartment package.

**BACKGROUND OF THE INVENTION**

15 Retinoids (e.g. retinol and retinyl esters) are common ingredients used in  
cosmetic products. Retinol (vitamin A) is an endogenous compound which occurs  
naturally in the human body and is essential for normal epithelial cell differentiation.  
Natural and synthetic vitamin A derivatives have been used extensively in the  
treatment of a variety of skin disorders and have been used as skin repair or renewal  
20 agents. Retinoic acid has been employed to treat a variety of skin conditions, e.g.,  
acne, wrinkles, psoriasis, age spots and discoloration. See e.g. Vahlquist, A. et al.,  
"Isotretinoin Treatment of Severe Acne Affects the Endogenous Concentration of  
Vitamin A in Sebaceous Glands," J. Invest. Dermatol., Vol. 94, pp. 496-498 (1990),  
Ellis, C.N. et. Al., "Pharmacology of Retinols in Skin," Basel, Karger, Vol. 3, (1989),  
25 pp. 249-252; and PCT Patent Application No. WO 93/19743.

**J6673(C)**  
**Y2-R566-EDG**

Retinoid metabolism, however may result in conversion of the retinoid to non-beneficial by-products, thus yielding a lesser amount of beneficial retinoic acid to treat skin conditions. Several prior art references, therefore, teach the use of a variety of natural actives for aiding in the treatment of skin conditions such as acne, wrinkles, psoriasis, age spots, and discoloration. For example, phytoestrogens (i.e., natural compounds which have estrogen-like activity and which are found in plants) have been increasingly used for cosmetic and therapeutic purposes. Estrogens and synthetic compounds which act like estrogens are known to increase the thickness of the dermal layer and reduce the wrinkle formation in the aging skin. Changes in the skin such as skin dryness, loss of skin elasticity and plumpness occurring after menopause are attributed to the lack of estrogen production. Estrogen therapy prevents or slows down many of the changes associated with aging skin (4) (Creidi et al., "Effect of a Conjugated Oestrogen Cream (Premarin®) on Aging Facial Skin," Maturitas, 19, p. 211-213, 1994).

Several phytoestrogens have been disclosed in the prior art for cosmetic benefits. For example, U.S. Patent No. 5,728,726 teaches the use of genistein for tyrosine kinase inhibitory activity. U.S. Patent Nos. 5,847,003 and 5,834,513 assigned to Avon disclose the use of oxacids and oxadiacids in combination with retinoids. Both Avon patents disclose the use of antioxidant bioflavonoids, such as genistein and daidzein, as optional ingredients.

It has been discovered, however, that phytoestrogens induce oxidation of retinol, and therefore contribute to retinol degradation. Although multi-compartment systems for delivering compositions have been described in the prior art, none disclose the need to separate phytoestrogens from retinoids. For example, U.S. Patent No. 5,914,116 issued to the assignee of the present invention describes two

**J6673(C)**  
**Y2-R566-EDG**

separate containers for separating two different skin actives to provide dual skin benefits with one compartment containing retinoids and the second compartment containing a second active providing a second and different benefit.

5           Therefore, there still exists a need for compositions that provide the skin benefits of retinoids along with the retinoid enhancing benefits of phytoestrogens.

**SUMMARY OF THE INVENTION**

10           A stable skin care product containing:

          a first composition comprising about 0.001% to about 10% of a retinoid;

15           a second composition comprising about 0.0001% to about 50% of at least one retinoid booster and about 0.001% to about 10% of a phytoestrogen;

          a first compartment for storing the first composition; and

20           a second compartment for storing the second composition, the first and second compartments being joined together.

**DETAILED DESCRIPTION**

25           As used herein, the term "comprising" means including, made up of, composed of, consisting and/or consisting essentially of. Except in the operating and comparative examples, or where otherwise explicitly indicated, all numbers in

**J6673(C)**  
**Y2-R566-EDG**

this description indicating amounts or ratios of materials or conditions of reaction, physical properties of materials and/or use are to be understood as modified by the word "about".

5           The inventive compositions contain, as a preferred ingredient, a retinoid, which is selected from retinyl esters, retinol, retinal and retinoic acid, preferably retinol or retinyl ester. The term "retinol" includes the following isomers of retinol: all-trans-retinol, 13-cis-retinol, 11-cis-retinol, 9-cis-retinol, 3,4-didehydro-retinol, 3,4-didehydro-13-cis-retinol; 3,4-didehydro-11-cis-retinol; 3,4-didehydro-9-cis-retinol. Preferred  
10 isomers are all-trans-retinol, 13-cis-retinol, 3,4-didehydro-retinol, 9-cis-retinol. Most preferred is all-trans-retinol, due to its wide commercial availability.

Retinyl ester is an ester of retinol. The term "retinol" has been defined above. Retinyl esters suitable for use in the present invention are C<sub>1</sub>-C<sub>30</sub> esters of retinol, preferably C<sub>2</sub>-C<sub>20</sub> esters, and most preferably C<sub>2</sub>, C<sub>3</sub>, and C<sub>16</sub> esters because they  
15 are more commonly available. Examples of retinyl esters include but are not limited to: retinyl palmitate, retinyl formate, retinyl acetate, retinyl propionate, retinyl butyrate, retinyl valerate, retinyl isovalerate, retinyl hexanoate, retinyl heptanoate, retinyl octanoate, retinyl nonanoate, retinyl decanoate, retinyl undecanoate, retinyl laurate, retinyl tridecanoate, retinyl myristate, retinyl pentadecanoate, retinyl heptadecanoate,  
20 retinyl stearate, retinyl isostearate, retinyl nonadecanoate, retinyl arachidonate, retinyl behenate, retinyl linoleate, retinyl oleate.

The preferred ester for use in the present invention is selected from retinyl  
25 palmitate, retinyl acetate and retinyl propionate, because these are the most commercially available and therefore the cheapest. Retinyl linoleate and retinyl oleate are also preferred due to their efficacy.

J6673(C)  
Y2-R566-EDG


Retinol or retinyl ester is employed in the inventive composition in an amount of about 0.001% to about 10%, preferably in an amount of about 0.01% to about 1%, most preferably in an amount of about 0.01% to about 0.5%.

5

It is believed that retinoids are enzymatically converted in the skin into retinoic acid according to the mechanism described in Chart 1.

10

## Retinol metabolism in the epidermis: enzyme names



ARAT/LRAT = Acyl Coenzyme A (CoA): Retinol Acyl Transferase/Lecithin:Retinol Acyl Transferase

15 CRABPII = Cellular Retinoic Acid Binding Protein II

It has been discovered, surprisingly, that certain compounds inhibit ARAT/LRAT, retinal reductase, CRABP II and retinoic acid oxidation (the latter catalyzed by cytochrome P450 systems), whereas certain other compounds enhance retinol dehydrogenase. The compounds are collectively termed herein as "boosters" and are coded as groups B1 through B5, as can be seen in Chart 1 hereinabove. The boosters, alone or in combination with each other, potentiate the action of a retinoid by increasing the amount of retinol available for conversion to retinoic acid and inhibiting the degradation of retinoic acid. The boosters act in conjunction with a retinoid (e.g. retinol, retinyl ester, retinal, retinoic acid), the latter being present endogenously in the skin. The preferred compositions, however, include a retinoid in the composition, co-present with a booster, to optimize performance.

The present invention includes, in part, a second composition containing about 0.0001% to about 50%, preferably about 0.001% to about 10%, most preferably about 0.001% to about 5% by weight of the composition of at least one booster compound, wherein the compound, either alone or at a combined concentration of 10mM, inhibits transglutaminase [in an in vivo transglutaminase assay] to more than 50%, and a cosmetically acceptable vehicle.

The boosters included in the inventive compositions are selected from the group consisting of:

- (a) Two boosters, wherein both are selected from the same group consisting of B2; B3; B4;

**J6673(C)**  
**Y2-R566-EDG**

(b) binary combinations of boosters selected from the group consisting of:

B1/B2; B1/B3; B1/B4; B1/B5; B2/B3, B2/B4; B2/B5, B3/B4; B3/B5;  
B4/B5;

(c) ternary combinations of boosters selected from the group consisting of:

B1/B2/B3; B1/B2/B4; B1/B2/B5; B1/B3/B4; B1/B3/B5;  
B1/B4/B5; B2/B3/B4; B2/B3/B5; B2/B4/B5; B3/B4/B5;

(d) quaternary combinations of boosters selected from the group consisting of:

B1/B2/B3/B4; B1/B2/B3/B5; B1/B2/B4/B5; B1/B3/B4/B5; B2/B3/B4/B5;  
and

(e) a combination of five groups of boosters: B1/B2/B3/B4/B5.

The preferred compositions include at least one booster from the different groups (i.e., groups (b) through (e) above). However, any combination of boosters chosen from the different groups may also be employed in the inventive compositions for desired boosting effects.

The compounds included in the present invention as boosters are first selected based on the ability of such compounds to pass, at a certain concentration listed in Table A, an in-vitro Microsomal Assay for a specific enzyme as described below under sections 2.1 through 2.7. The compound (alone or in combination with another booster) is then subjected to an in vitro transglutaminase assay described below, at an individual or combined concentration of 10 mM. If such combination inhibits transglutaminase to more than 50%, then it is suitable for use in the present invention. If a booster was tested individually, and passes the transglutaminase assay, then it

**J6673(C)**  
**Y2-R566-EDG**

may be combined with another booster or combination that passes the transglutaminase assay.

Preferred compositions according to the present invention contain  
5 combinations of boosters which at an individual concentration of 10 mM inhibit transglutaminase to more than 50%.

The term "conditioning" as used herein means prevention and treatment of dry  
skin, acne, photodamaged skin, appearance of wrinkles, age spots, aged skin,  
10 increasing stratum corneum flexibility, lightening skin color, controlling sebum excretion and generally increasing the quality of skin. The composition may be used to improve skin desquamation and epidermal differentiation.

A booster is a compound which passes an in vitro Microsomal Assay described  
15 below in sections 2.1 through 2.7. A compound suitable for use in the present invention inhibits or enhances, at a concentration listed in Table A an enzyme, to at least a broad % listed in Table A.

20

25



J6673(C)  
Y2-R566-EDG

TABLE A

Booster Test Concentrations and % Inhibition/Increase

ARAT / LRAT Assay *(To identify B1 boosters)*

Invention	Compound Concentration	% Inhibition
Broad	100 µM	> 10%
Preferred	100 µM	> 25%
Most Preferred	100 µM	> 40%
Optimum	100 µM	> 50%

Retinol Dehydrogenase Assay *(To identify B2 boosters)*

Invention	Compound Concentration	% Increase
Broad	100 µM	> 10%
Preferred	100 µM	> 15%
Most Preferred	100 µM	> 20%
Optimum	100 µM	> 25%

5

Retinal Reductase Assay *(To identify B3 boosters)*

Invention	Compound Concentration	% Inhibition
Broad	100 µM	> 5%
Preferred	100 µM	> 10%
Most Preferred	100 µM	> 20%
Optimum	100 µM	> 35%

CRABPII Antagonist Assay *(To identify B4 boosters)*

Invention	Compound : RA Ratio	% Inhibition
Broad	7000 : 1	> 25%
Preferred	7000 : 1	> 50%
Most Preferred	70 : 1	> 25%
Optimum	70 : 1	> 50%

10

**J6673(C)**  
**Y2-R566-EDG**

**Retinoic Acid Oxidation Assay (To identify B5 boosters)**

Invention	Compound Concentration	% Inhibition
Broad	100 $\mu$ M	> 25%
Preferred	100 $\mu$ M	> 45%
Most Preferred	100 $\mu$ M	> 70%
Optimum	100 $\mu$ M	> 80%

The in vitro Microsomal Assays employed for determining the suitability of the inclusion of the compound in the inventive compositions are as follows:

5

**1. Materials**

All-trans-retinol, all-trans-retinoic acid, palmitoyl-CoA, dilauroyl phosphatidyl choline, NAD, and NADPH were purchased from Sigma Chemical Company. Stock solutions of retinoids for the microsomal assays were made up in HPLC grade acetonitrile. All retinoid standard stock solutions for HPLC analysis were prepared in ethanol, stored under atmosphere of N<sub>2</sub> at -70°C and maintained on ice under amber lighting when out of storage. Other chemicals and the inhibitors were commercially available from cosmetic material suppliers or chemical companies such as Aldrich or International Flavors and Fragrances.

15

**2. Methods**

2.1 Isolation of RPE microsomes (modified from J.C. Saari & D.L. Bredberg, "CoA and Non-CoA Dependent Retinol Esterification in Retinal Pigment Epithelium", J. Biol. Chem. 263, 8084-8090 (1988)).

20

50 frozen hemisected bovine eyecups, with the retina and aqueous humor removed were obtained from W. L. Lawson Co., Lincoln, NE, USA. The eyes were thawed

**J6673(C)**  
**Y2-R566-EDG**

overnight and the colored iridescent membrane was removed by peeling with forceps. Each eyecup was washed with 2x 0.5mL cold buffer (0.1M PO<sub>4</sub> / 1mM DTT / 0.25M sucrose, pH 7) by rubbing the darkly pigmented cells with an artist's brush or a rubber policeman. The cell suspension was added to the iridescent membranes and the suspension was stirred for several minutes in a beaker with a Teflon stir bar. The suspension was filtered through a coarse filter (Spectra / Por 925μ pore size polyethylene mesh) to remove large particles, and the resulting darkly colored suspension was homogenized using a Glas-Col with a motor driven Teflon homogenizer. The cell homogenate was centrifuged for 30 min. at 20,000g (Sorvaal model RC-5B centrifuge with an SS34 rotor in 2.5x10cm tubes at 14,000 RPM). The resulting supernatant was subjected to further centrifugation for 60 min. at 150,000g (Beckman model L80 Ultracentrifuge with an SW50.1 rotor in 13x51mm tubes at 40,000 RPM). The resulting pellets were dispersed into ~5mL 0.1M PO<sub>4</sub> / 5mM DTT, pH 7 buffer using a Heat Systems Ultrasonics, Inc. model W185D Sonifier Cell Disruptor, and the resulting microsomal dispersion was aliquoted into small tubes and stored at -70°C. The protein concentrations of the microsomes were determined using the BioRad Dye binding assay, using BSA as a standard.

2.2 Isolation of rat liver microsomes (modified from R. Martini & M. Murray, "Participation of P450 3A Enzymes in Rat Hepatic Microsomal Renitoic Acid 4-Hydroxylation", Archives Biochem. Biophys. 303, 57-66 (1993)).

Approximately 6 grams of frozen rat liver (obtained from Harlan Sprague Dawley rats from Accurate Chemical and Scientific Corp.) were homogenized in 3 volumes of 0.1M tris / 0.1M KCl / 1mM EDTA / 0.25M sucrose, pH 7.4 buffer using a Brinkmann Polytron. The resulting tissue suspension was further homogenized in the motor driven Teflon homogenizer described above. The resulting homogenate was

**J6673(C)**  
**Y2-R566-EDG**

successively centrifuged for 30 min. at 10,000g, 30 min. at 20,000g, and 15 min. at 30,000g, and the resulting supernatant was ultracentrifuged for 80 min. at 105,000g. The pellet was sonicated in ~5mL of 0.1M PO<sub>4</sub> / 0.1mM EDTA / 5mM MgCl<sub>2</sub> , pH 7.4 buffer as described above and stored as aliquots at -70°C. Protein concentrations  
5 were determined as described above.

**2.3 Assay for ARAT and LRAT activity (To identify B1)**

The procedure below is a modification of a method described in J.C. Saari & D.L. Bredberg, "ARAT & LRAT Activities of Bovine Retinal Pigment Epithelial Microsomes",  
10 Methods Enzymol. 190, 156-163 (1990). The following buffer was prepared and stored at 4°C: 0.1M PO<sub>4</sub> / 5mM dithiothreitol, pH 7.0 (PO<sub>4</sub> / DTT). On the day of the assay, add 2mg BSA per mL of buffer to give a PO<sub>4</sub> / DTT / BSA working buffer. 1mM retinol substrate was prepared in acetonitrile and stored in amber bottles under nitrogen gas at -20°C. Solutions of 4mM Palmitoyl-CoA in working buffer (stored in aliquots) and  
15 4mM dilauroyl phosphatidyl choline in ethanol were prepared and stored at -20°C. Inhibitors were prepared as 10mM stock solutions in H<sub>2</sub>O, ethanol, acetonitrile or DMSO. The quench solution was prepared using pure ethanol containing 50µg/mL butylated hydroxytoluene (BHT), and a hexane solution containing 50µg/mL BHT was used for the extractions.

20

To a 2 dram glass vial, add the following in order: PO<sub>4</sub> / DTT / BSA buffer to give a total volume of 500µL, 5µL acyl donor (4mM palmitoyl-CoA and/or dilauroyl phosphatidyl choline), 5µL inhibitor or solvent blank (10mM stock or further dilutions) followed by approximately 15µg of RPE microsomal protein (approximately 15µL of a  
25 ~1mg/mL microsomal protein aliquot). Incubate for 5 min. at 37°C to equilibrate the reaction temperature and then add 5µL 1mM retinol. Cap the vials, vortex for 5 seconds and incubate for 30-90 minutes at 37°C. Quench the reaction by adding

**J6673(C)**  
**Y2-R566-EDG**

0.5mL ethanol / BHT. Extract the retinoids by adding 3mL hexane / BHT, vortex the tubes for several seconds several times and centrifuge the tubes at low speed for 5 min. to quickly separate the layers. Remove the upper hexane layer into a clean vial, and re-extract the aqueous layer with another 3mL hexane / BHT, as described  
5 above. Combine the hexane layers and evaporate the hexane by drying at 37°C under a stream of nitrogen gas on a heated aluminum block. Store the dried residue at -20°C until HPLC analysis. Quantitate the amount of retinyl palmitate and retinyl laurate for ARAT and LRAT activity, respectively, by integration of the HPLC signal as described below.

10 Note that the incubation solution contains 40µM acyl donor, 100µM or less inhibitor, 10µM retinol, approximately 30µg/mL microsomal protein, and nearly 0.1M PO<sub>4</sub>, pH 7 / 5mM DTT / 2mg/mL BSA. All steps subsequent to the addition of retinol were done in the dark or under amber lights.

15 **2.4 Assay for Retinol Dehydrogenase Activity** (To identify B2)

The following stock solutions were prepared:

20 50mM KH<sub>2</sub>PO<sub>4</sub>, pH 7.4 buffer, sterile filtered.

10mM all trans Retinol (Sigma R7632) in DMSO.

200mM Nicotinamide adenine dinucleotide phosphate, sodium salt (NADP) (Sigma N0505) in sterile water.

40mM test compound in appropriate solvent (water, buffer, ethanol, chloroform or  
25 DMSO).

1:10 dilution of rat liver Microsomes in 50mM KH<sub>2</sub>PO<sub>4</sub>, pH 7.4 buffer (4ug/ul).

**J6673(C)**  
**Y2-R566-EDG**

In a two-dram glass vial with screw cap, add the following in order:

Buffer to give a final volume of 400 $\mu$ l

- 25 $\mu$ l diluted Microsomes (final = 100ug) – use boiled Microsomes for controls and  
5 regular Microsomes for test samples.

4 $\mu$ l of 200mM NADP (final = 2mM)

1 $\mu$ l of 40mM test compound (final = 100 $\mu$ M)

8 $\mu$ l of 10mM retinol (final = 200 $\mu$ M)

- 10 Incubate vials in a 37°C shaking water bath for 45 minutes. Add 500 $\mu$ l ice-cold ethanol to each vial to quench the reaction. Extract the retinoids twice with ice cold hexane (2.7ml per extraction). Retinyl acetate (5 $\mu$ l of a 900 $\mu$ M stock) is added to each vial during the first extraction as a means of monitoring the extraction efficiency in each sample. Samples were vortexed for ten seconds before gently centrifuging for  
15 five minutes at 1000rpm, 5°C in a Beckman GS-6R centrifuge. The top hexane layer containing the retinoids is removed from the aqueous layer after each extraction to a clean two-dram vial. Evaporate off the hexane under a gentle stream of nitrogen gas. Store the dried residue at –20°C until HPLC analysis.

- 20 2.5 Assay for Retinal Reductase Activity (To identify B3)

All stock solution were prepared as above with the following substitutions:

- 10mM all trans Retinaldehyde (Sigma R2500) in DMSO – instead of retinol.  
25 200mM, Nicotinamide adenine dinucleotide phosphate, reduced form, tetrasodium salt (NADPH) (Sigma N7505) in sterile water – instead of NADP.

**J6673(C)**  
**Y2-R566-EDG**

In a two-dram glass vial with screw cap, add the following in order:

Buffer to give a final volume of 400 $\mu$ l

- 25 $\mu$ l diluted Microsomes (final = 100 $\mu$ g) – use boiled Microsomes for controls and  
5 regular Microsomes for test samples.

4 $\mu$ l of 200mM NADPH (final = 2mM)

1 $\mu$ l of 40mM test compound (final = 100 $\mu$ M)

3 $\mu$ l of 10mM retinaldehyde (final = 75 $\mu$ M)

- 10 Follow the same incubation and extraction procedure as detailed above.

**2.6 Assay for CRABP II antagonists (To identify B4)**

**2.6.1 Synthesis of CRABP II**

- 15 a. System of expression

The gene CRABP II was cloned in pET 29a-c(+) plasmid (Novagen). The cloned gene was under control of strong bacteriophage T7 transcription and translation signals. The source of T7 polymerase was provided by the host cell E.coli BLR(DE3)pLysS (Novagen). The latter has a chromosomal copy of T7 polymerase under lacUV5  
20 control, induced by the presence of IPTG. The plasmid was transferred into E. coli BLR(DE3)pLysS cells by transformation according to the manufacturer protocol (Novagen).

**b. Induction**

- 25 An overnight culture of the transformed cells was diluted 1:100 into 2xYT containing 50  $\mu$ g/mL kanamycin and 25 $\mu$ g/mL chloramphenicol. The cells grew while shaking at 37°C until the OD at 600 nm reached 0.6-0.8. Then IPTG was added to a final

**J6673(C)**  
**Y2-R566-EDG**

concentration of 1mM and the culture was incubated for an additional two hours. The cells were harvested by centrifugation at 5000g for 10 minutes at room temperature. The pellet was stored at -20°C.

5    2.6.2 Purification

Purification was performed according to the method described in A. W. Norris and E. Li, "Generation and characterization of cellular retinoic acid-binding proteins from *Escherichia coli* expression systems. *Methods Enzymol*, 1997;282:3-13.

a. Lysis

10    The frozen pellet was thawed at RT and resuspended in 1-2 pellet volumes of freshly prepared lysis buffer (50 mM Tris-HCl, pH 8, 10%(w/v) sucrose, 1 mM EDTA, 0.05%(w/v) sodium azide, 0.5 mM DTT, 10 mM MnCl<sub>2</sub>, 2.5 mM phenylmethylsulfonyl fluoride, 2.5 mM benzamidine, 6µg/mL DNase). The lysate was incubated for 30 min at room temperature. Further lysis was accomplished by sonication (six 30-sec bursts at 10,000 psi alternated with five 30-sec delay on ice). The insoluble fraction of the  
15    lysate was removed by centrifugation at 15000 rpm 1 hour at 4°C and the supernatant is stored at -20°C.

b. Gel filtration on Sephacryl S300

20    The supernatant from step a. was loaded onto a 2.5x100 cm column of sephacryl S-300 (Pharmacia) at room temperature. The elution buffer was 20 mM Tris-HCl, pH 8, 0.5mM DTT, 0.05% sodium azide (buffer A). The flow rate was 2mL/min. Collected 2-mL fractions were checked for ultraviolet absorbance at 280 nm. The fractions representing the peaks were examined by SDS-page for the presence of CRABP II .

25

c. Anion-exchange chromatography



**J6673(C)**  
**Y2-R566-EDG**

2 mL of gel filtration fractions containing CRABP II were loaded onto a quaternary amine anion-exchange column FPLC (Fast Protein Liquid Chromatography) type monoQ (Pharmacia). CRABP II was eluted using a gradient buffer from 100% buffer A to 30% buffer B (100 % buffer B = buffer A + 250 mM NaCl) over a 20-min period at room temperature. 1 mL-fractions were collected every minute. Once more, the presence of CRABP II was checked by SDS page. CRABP II was stored at 4°C before freeze-drying using a Micromodulyo 1.5K with vial platform attachment (Edwards High Vacuum International). The desiccated samples were stored at room temperature until their use in the binding assay.

d. Detection of the presence of CRABP II

The expression and purification of CRABP II was validated using denaturing SDS-polyacrylamide gel electrophoresis (SDS-PAGE) analysis on a 7-15% polyacrylamide gel (Biorad). 10 µL samples were mixed with 10 µL of 2X loading buffer (100 mM Tris-HCl pH6.8, 4% SDS, 0.2% BPB, 20% glycerol, 1mM DTT) and denatured by heating (2 min at 80°C). The samples were loaded onto the gel that was immersed in a 1X Tris-glycine buffer (Biorad) and a constant current (25 mA) was applied for 1 hour at room temperature. After Coomassie blue staining, the protein was identified according to its molecular weight as determined with the Benchmark prestained protein ladder (Gibco BRL).

A western blot was used to confirm the presence of CRABP II. The proteins separated on the SDS-PAGE were transferred on an Immobilon-P transfer membrane (Millipore) using a Biorad cassette. The transfer occurred in 1X Tris-glycine buffer (Biorad) + 10% methanol. An electrical current (60 mA) was applied for 3 hours to allow the protein to migrate through the membrane. Afterwards, the membrane was blocked with 5% dry milk in 1X TBS for one hour at room temperature and probed with primary antibodies

**J6673(C)**  
**Y2-R566-EDG**

to CRABP II (1/1000 dilution of mouse anticonal 5-CRA-B3) in the same buffer at 4°C overnight. The following day, the membrane was washed with PBS (3 x 5 minutes) and then incubated with 1:2000 dilution of the secondary antibody, peroxidase conjugated anti-mouse antibody (ECLTM, Amersham), for 1 hour at room temperature. The membrane was washed with 1xPBS (3x 5 minutes) and the protein was detected using ECL detection kit according to the manufacturer instruction (Amersham).

The concentration of purified CRABP II was determined using BSA kit (Pierce).

10    2.6.3 Radioactive Binding assay

220 pmol of CRABP II was incubated in 20 mM Tris-HCl buffer pH 7.4 with 15 pmol of radioactive all trans retinoic acid (NEN) in a total volume of 70µL. For the competitive assay, another ligand in excess (6670:1, 670:1 or 70:1) was added to the mix. The reaction occurred for one hour at room temperature in the dark. In order to separate the unbound all-trans retinoic acid from the bound all-trans retinoic acid, a 6kD cut-off minichromatography column (Biorad) was used. The storage buffer was discarded using a Microplex manifold for according to the manufacturer instruction (Pharmacia). The samples were loaded onto the column and the separation occurred by gravity over a 30-min period. Retinoic acid ("RA") bound to CRABP II appeared in the filtrate while free RA remained in the column. The radioactivity of the filtrate was measured by scintillation counter.

25    2.7 Assay for NADPH dependent retinoic acid oxidation (To identify B5)

The procedure below is a modification of a method described in R. Martini & M. Murray, "Participation of P450 3A Enzymes in Rat Hepatic Microsomal Retinoic Acid

**J6673(C)**  
**Y2-R566-EDG**

4-Hydroxylation", Archives Biochem. Biophys. 303, 57-66 (1993). Prepare the following assay buffer and store at 4°C: 0.1M PO<sub>4</sub> / 0.1mM EDTA / 5mM MgCl<sub>2</sub>, pH 7.4. On the day of the assay, prepare a 60mM NADPH solution in buffer. Prepare inhibitor stocks, acidified ethanol / BHT quench solution, and hexane / BHT as described above. A working 1mM retinoic acid solution was prepared by dilution of a 15mM stock (in DMSO) with ethanol.

To a 2 dram vial, add the following in order: assay buffer to give a final volume of 500µL, 20µL 60mM NADPH, 5µL inhibitor or solvent blank, followed by approximately 2mg of rat liver microsomal protein.

Incubate for 5 min. at 37°C, then add 5µL working 1mM retinoic acid solution. Continue incubation for 60min. at 37°C - do not cap the vials, since the oxidation process requires molecular O<sub>2</sub> in addition to NADPH. Quench with acidified ethanol / BHT and extract with hexane / BHT as described above. Quantitate the quickly eluting polar retinoic acid metabolites (presumed to be 4-oxo retinoic acid) by integration of the HPLC signal, as described below.

Note that all steps subsequent to the addition of retinoic acid were done in the dark or under amber lights. The final incubation solution contains 2.4mM NADPH, 100µM or less inhibitor, 10µM retinoic acid, approximately 4mg/mL rat liver microsomal protein and nearly 0.1M PO<sub>4</sub> / 0.1mM EDTA / 5mM MgCl<sub>2</sub>.

**HPLC analysis of individual retinoids**

Samples for retinoid quantitation by HPLC were prepared by dissolving the residue in each vial with 100µL of methanol. The solution was transferred to a 150µL glass conical tube within a 1mL shell vial, capped tightly, and placed inside a Waters 715

**J6673(C)**  
**Y2-R566-EDG**

Autosampler. Aliquots of 60µL were injected immediately and analyzed for retinoid content.

5 The chromatography instrumentation consisted of a Waters 600 gradient controller / pump, a Waters 996 Photodiode Array detector and a Waters 474 Scanning Fluorescence detector. Two HPLC protocols were used for retinoid analysis. For the ARAT and LRAT assay, the separation of retinol and retinol esters was performed with a Waters 3.9x300mm C18 Novapak reverse-phase analytical column and Waters Sentry NovaPak C18 guard column with an 80:20(v/v) methanol / THF isocratic mobile  
10 phase adjusted to a flow rate of 1mL/min. for 10 min. The eluate was monitored for absorbance at 325nm and fluorescence at 325ex/480em. A shorter Waters 3.9x150mm C18 Novapak reverse-phase analytical column and Waters Sentry NovaPak C18 guard column were used to separate retinoid acids and alcohols for the retinol and retinoic acid oxidation assays utilizing a modification of a gradient system  
15 described by A. B. Barua, "Analysis of Water-Soluble Compounds: Glucoronides", Methods Enzymol. 189, 136-145 (1990). This system consisted of a 20 min. linear gradient from 68:32(v/v) methanol/ water containing 10mM ammonium acetate to 4:1(v/v) methanol:dichloromethane followed by a 5 min. hold at a flow rate of 1mL/min.. The column eluate was monitored from 300nm to 400nm.

20

These protocols were selected based on their ability to clearly resolve pertinent retinoid acids, alcohols, aldehydes, and/or esters for each assay and relative quickness of separation. Identification of individual retinoids by HPLC was based on an exact match of the retention time of unknown peaks with that of available authentic retinoid  
25 standards and UV spectra analysis (300-400nm) of unknown peaks against available authentic retinoids.

**J6673(C)**  
**Y2-R566-EDG**

The boosters suitable for further testing in the transglutaminase assay include but are not limited to the boosters listed in Tables B1 through B5 below.

2025-04-04 14:00:00

**J6673(C)**  
**Y2-R566-EDG**

**ARAT/LRAT Inhibitors (B1)**

Class	Compound	% Inhibition Overall TG (- ROH/RE)	Overall TG (IC 50)	% Inhibition ARAT (10µm)	% Inhibition ARAT(100µm)	% Inhibition LRAT (10µm)	% Inhibition LRAT (100µm)
Carotenoid	Croctin		3.75E-05	15%	34%	0	15%
Fatty Acid Amides & Other Surfactants	Acetyl Sphingosine		6.78E-06	19%+/- 12	62% +/- 11	10% +/- 10	50% +/- 18
Fatty Acid Amides & Other Surfactants	C13 Beta-Hydroxy Acid/Amide	17%			28%		25%
Fatty Acid Amides & Other Surfactants	Castor Oil MEA		3.25E-05				
Fatty Acid Amides & Other Surfactants	Cocamidopropyl Betaine				25%		
Fatty Acid Amides & Other Surfactants	Coco Hydroxyethylimidazoline		2.84E-07		68%		68%
Fatty Acid Amides & Other Surfactants	Cocoamide-MEA (or Cocoyl Monoethanolamide)	11%			13%		34%
Fatty Acid Amides & Other Surfactants	Glycerol-PCA-Oleate				41% +/- 6		58% +/- 2
Fatty Acid Amides & Other Surfactants	Hexanoamide				20%		
Fatty Acid Amides & Other Surfactants	Hexanoyl Sphingosine		9.99E-05		28% +/-4		37% +/-9
Fatty Acid Amides & Other Surfactants	Hydroxyethyl-2-Hydroxy-C12 Amide		3.29E-05		35%		35%
Fatty Acid Amides & Other Surfactants	Hydroxyethyl-2-Hydroxy-C16 Amide				25%		30%
Fatty Acid Amides & Other Surfactants	Lauroyl Sarcosine				20%		
Fatty Acid Amides & Other Surfactants	Lidocaine				12%		0
Fatty Acid Amides & Other Surfactants	Linoleamide-DEA (or Linoleoyl Diethanolamide)	59%		12% +/-13	43% +/-3	11% +/-9	51% +/- 15
Fatty Acid Amides & Other Surfactants	Linoleamide-MEA (or Linoleoyl Monoethanolamide)		1.61E-05	14%	35%	20% +/- 8	35%
Fatty Acid Amides & Other Surfactants	Linoleamidopropyl Dimethylamine				69% +/-18		75% +/-4
Fatty Acid Amides & Other Surfactants	Melinamide				64% +/-15		43% +/-2 21
Fatty Acid Amides & Other Surfactants	Myristoyl Sarcosine				41% +/- 14		11% +/-11

**J6673(C)  
Y2-R566-EDG**

Class	Compound	% Inhibition Overall TG (- ROH/RE)	Overall TG (IC 50)	% Inhibition ARAT (10µm)	% Inhibition ARAT(100µm)	% Inhibition LRAT (10µm)	% Inhibition LRAT (100µm)
Fatty Acid Amides & Other Surfactants	Oleyl Betaine		2.80E-05	47%			
Fatty Acid Amides & Other Surfactants	Palmitamide-MEA			6%	23%	12%	33%
Fatty Acid Amides & Other Surfactants	Stearylhydroxyamide				10%		10%
Fatty Acid Amides & Other Surfactants	Utrecht-1	21%		43%	54%	51%	48% +/- 6
Fatty Acid Amides & Other Surfactants	Utrecht-2		3.47E-06	42%	83% +/-9	51%	92% +/-3
Flavonoids	Naringenin				33%		14%
Fragrances	Allyl Alpha-Ionone			16% +/-14	22% +/-23	17% +/- 10	36%/- 7
Fragrances	Alpha-Damascone		3.35E-04	67% +/- 27	83% +/- 12	87% +/- 6	98% +/- 1
Fragrances	Alpha=Ionone		9.27E-04		45% +/- 27		49% +/- 30
Fragrances	Alpha-Methyl Ionone				67%		77%
Fragrances	Alpha-Terpineol				26%		25%
Fragrances	Beta-Damascone			45%	84%	52%	92%
Fragrances	Brahmanol				70%		75%
Fragrances	Damascenone			23%	70%	29%	79%
Fragrances	Delta-Damascone			58%	87%	64%	95%
Fragrances	Dihydro Alpha-Ionone				13%		18%
Fragrances	Ethyl Safranate				51%		49%
Fragrances	Fenchyl Alcohol				12%		4%
Fragrances	Gamma-Methyl Ionone				21%		38%
Fragrances	Isobutyl Ionone				8%		45%
Fragrances	Isocyclogeraniol				18%		16%
Fragrances	Isodamascone				80%		92%

**J6673(C)  
Y2-R566-EDG**

Class	Compound	% Inhibition Overall TG (- ROH/RE)	Overall TG (IC 50)	% Inhibition ARAT (10µm)	% Inhibition ARAT(100µm)	% Inhibition LRAT (10µm)	% Inhibition LRAT (100µm)
Fragrances	Lylal		1.27E-04	76%	71%		
Fragrances	Santalone			23%			12%
Fragrances	Santanol			15%			43%
Fragrances	Timberol			34%			33%
Fragrances	Tonalid			50%			33%
Fragrances	Traseolide			41%			21%
Miscellaneous	Coco Trimethylammonium Cl-			27%			
Miscellaneous	Urosolic Acid		1.46E-06	21%			28%
Noncyclic Fragrances	Citral			20%			
Noncyclic Fragrances	Citronellol			30%			0
Noncyclic Fragrances	Farnesol		9.35E-05	23%+/- 18	53% +/- 18	10% +/- 7	53% +/- 19
Noncyclic Fragrances	Geraniol		7.83E-03	13%	32%		
Noncyclic Fragrances	Geranyl Geraniol			38% +/- 12	81% +/- 6	16% +/- 9	77% +/- 13
Noncyclic Fragrances	Linatool				28%		0
Noncyclic Fragrances	Nonadieneal				20%		
Noncyclic Fragrances	Pseudoionone				12%		37%
Phospholipid	Diocetylphosphatidyl Ethanolamine						
Urea	Dimethyl Imidazolidinone	22%		23%	50% +/- 2	0	17% +/- 17
Urea	Imidazolidinyl Urea	35%					



J6673(C)  
Y2-R566-EDG

### Retinol Dehydrogenase Activators (B2)

Class	Compound	% Increase Retinol Dehydrogenase
Phospholipid	Phosphatidyl Choline	21% increase
Phospholipid	Sphingomyelin	26% increase

### Retinaldehyde Reductase Inhibitors (B3)

Class	Compound	Overall TG (IC 50)	% Inhibition Retinal Reductase
Aldehyde	Vanillin	9.70E-03	6%
Fatty Acid	Arachidic Acid		20%
Fatty Acid	Arachidic Acid		49%
Fatty Acid	Linoleic Acid	1.63E-04	62% +/-2
Fatty Acid	Linolenic Acid	1.34E-04	54% +/-16
Fatty Acid	Myristic Acid	1.72E-05	26%
Miscellaneous	Amsacrine	6.26E-06	22% + /-8
Miscellaneous	Carbenoxolone	3.61E-07	26% +/-2
Miscellaneous	Glycyrrhetic Acid	8.64E-06	38% +/- 1
Phospholipid	Phosphatidyl ethanolamine		37%

J6673(C)  
Y2-R566-EDG

CRABPII Antagonists (B4)

Class	Compound	Overall TG (IC 50)	% Inhibition CRABPII
Fatty Acid	Elaidic Acid	6.50E-05	>50%
Fatty Acid	Hexadecanedioic Acid	1.30E-04	>50%
Fatty Acid	12-Hydroxystearic Acid	2.91E-05	>50%
Fatty Acid	Isostearic Acid	6.88E-05	>50%
Fatty Acids	Linseed Oil		>50%

5

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100

J6673(C)  
Y2-R566-EDG

Retinoic Acid Oxidation Inhibitors (B5)

Class	Compound	Overall TG (IC 50)	% Inhibition Retinoic Acid (10µM)	% Inhibition Retinoic Acid (100µM)
Imidazole	Bifonazole		89%	100%
Imidazole	Climbazole	4.47E-06	80%	92%
Imidazole	Clotrimazole		76%	85%
Imidazole	Econazole		88%	100%
Imidazole	Ketoconazole	1.85E-07	84%	84%
Imidazole	Miconazole	2.78E-07	74%	86%
Fatty Acid Amides & Other Sufactants	Lauryl Hydroxyethylimidazoline	4.67E-07		
Fatty Acid Amides & Other Sufactants	Oleyl Hydroxyethylimidazoline	3.02E-05	54%	80%
Flavanoids	Quercetin	6.29E-05	40%	74%
Coumarin	Coumarin			
Quinoline	(7H-Benzimidazo[2, 1-a]Benz[de]-Isoquinolin-7-one	8.59E-07		
Quinoline	Hydroxyquinoline (Carbostyryl)	3.64E-04		
Quinoline	Metrapone (2-Methyl-1, 2-di-3-Pyridyl-1-Propane			47%

**J6673(C)**  
**Y2-R566-EDG**

The boosters or combinations thereof inhibit transglutaminase (hereinafter "Tgase") in a transglutaminase assay described below to at least 50% at a concentration of 10mM.

5

TGase Assay		
Invention	Compound Concentration	% Inhibition
Broad	10 mM	> 50%
Preferred	1 mM	> 50%
Most Preferred	100 $\mu$ M	> 50%
Optimum	10 $\mu$ M	> 50%

**Transglutaminase Assay and Keratinocyte Differentiation**

10 During the process of terminal differentiation in the epidermis, a 15nm thick layer of protein, known as the cornified envelope (CE) is formed on the inner surface of the cell periphery. The CE is composed of numerous distinct proteins which have been cross-linked together by the formation of N<sup>ε</sup>-( $\gamma$ -glutamyl) lysine isodipeptide bonds catalyzed by the action of at least two different transglutaminases

15 (TGases) expressed in the epidermis. TGase I is expressed in abundance in the differentiated layers of the epidermis, especially the granular layer, but is absent in the undifferentiated basal epidermis. Thus TGase I is a useful marker of epidermal keratinocyte differentiation with high TGase I levels indicating a more differentiated state. An ELISA based TGase I assay, using a TGase I antibody, was used to assess

20 the state of differentiation of the cultured keratinocytes in the examples that follow.

Keratinocytes (cultured as described above) were plated in 96 well plates at a density of 4,000-5,000 cells per well in 200 $\mu$ l media. After incubation for

**J6673(C)**  
**Y2-R566-EDG**

two to three days, or until cells are ~50% confluent, the media was changed to media containing test compounds (five replicates per test). The cells were cultured for a further 96 hours after which time the media was aspirated and the plates stored at -70°C. Plates were removed from the freezer, and the cells were washed twice  
5 with 200µl of 1x PBS. The cells were incubated for one hour at room temperature (R/T) with TBS/5% BSA (wash buffer, bovine serum albumin). Next the TGase primary antibody was added: 50µl of monoclonal anti-Tgase I Ab B.C. diluted 1:2000 in wash buffer. The primary antibody was incubated for 2 hours at 37°C and then rinsed 6x with wash buffer. Cells were then incubated with 50µl of secondary  
10 antibody (Fab fragment, peroxidase conjugated anti-mouse IgG obtaining from Amersham) diluted 1:4,000 in wash buffer for two hours at 37°C, then rinsed three times with wash buffer. Following the rinse with washing buffer, the cells were rinsed 3x with PBS. For colourimetric development, the cells were incubated with  
15 100µl substrate solution (4 mg o-phenylenediamine and 3.3 µl 30% H<sub>2</sub>O<sub>2</sub> in 10ml 0.1M citrate buffer pH 5.0) for exactly five minutes, R/T, in darkness (under aluminum foil). The reaction was stopped by the addition of 50µl 4N H<sub>2</sub>SO<sub>4</sub>. The absorbance of samples was read at 492nm in a 96 well plate UV spectrophotometer. Out of the five replicates, four were treated with both antibodies, the fifth one was  
20 use as a Tgase background control. TGase levels were determined and expressed as percentage control.

Transglutaminase levels were determined and expressed in the Tables B1 through B5 above either as:

(i) % (booster + retinol inhibition / control inhibition) - % (ROH inhibition / control inhibition), which measures the added effect of booster + retinol induced TGase inhibition over retinol alone, or

**J6673(C)  
Y2-R566-EDG**

(ii) as an IC<sub>50</sub> value when the inhibitory effect of multiple booster concentrations was examined - this provides the concentration of booster which, in combination with a constant retinol concentration of 10<sup>-7</sup>M, inhibits TGase by 50%.

5

It is the IC<sub>50</sub> value that is used as a benchmark in the present invention.

**Best Groups of Boosters for testing in Transglutaminase assay**

10

**B1 Compounds**

1. Fatty Acid Amides	These are readily commercially available and have the added advantage of being surfactants and thus help generate emulsions suitable for cosmetic preparations.
2. Ceramides	These can additionally act as precursors of stratum corneum barrier ceramides.
3. Carotenoids	These can offer some UV protection and and act as natural colorants.
4. Flavanoids	Natural antioxidants.
5. Cyclic fragrances	These are readily commercially available and additionally can be used to fragrance the product.
6. Non-cyclic fragrances	These can be used to fragrance the product.
7. Phospholipid analogues	These can be utilised by skin cells to nourish the generation of barrier components.
8. Ureas	These are readily commercially available and can also act as preservatives for the product.

15 **B2 Compounds**

1. Phosphatidyl choline	Most preferred as most active activator of Retinol Dehydrogenase
2. Sphingomyelin	

**J6673(C)**  
**Y2-R566-EDG**

### B3 Compounds

Arachidonic Acid Linoleic Acid Linolenic Acid Myristic Acid	Fatty Acids which can be useful in maintaining stratum corneum barrier
Linoleic Acid Linolenic Acid	Essential Fatty Acids
Arachidonic Acid Myristic Acid	Non-essential fatty acids
Glycyrrhetic Acid	Polycyclic triterpene carboxylic acid which is readily obtained from plant sources.
Phosphatidyl ethanolamine	Can be incorporated into cellular membranes.

5

### B4 Compounds

Hexadecanedioic acid 12-hydroxystearic acid Isostearic acid	Saturated fatty acids.
Linseed oil Elaidic acid	Unsaturated fatty acids
Elaidic acid Isostearic acid Hexadecanedioic acid	Solid at room temperature
Linseed oil 12-hydroxystearic acid	Liquid at room temperature

10

J6673(C)  
Y2-R566-EDG

## B5 Compounds

Bifonazole Climbazole Clotrimazole Econazole Ketoconazole Miconazole	Antimicrotics
Climbazole	Readily commercially available
Lauryl hydroxyethylimidazoline	Compounds which are readily commercially available and have the added advantage of being surfactants and thus help generate emulsions suitable for cosmetic preparations.
Quercetin	Naturally occurring flavanoid which has antioxidant properties.
Coumarin	Natural colorant
Quinolines Isoquinolines	
Metyrapone	

## PHYTOESTROGEN

As part of the present invention, it has been surprisingly found that phytoestrogens synergistically improve the skin benefits of retinoids. Essentially, phytoestrogens increase the sensitivity of the skin to retinoids.

Therefore, the present invention contains about 0.001% to about 10% of at least one phytoestrogen in the second composition.

Phytoestrogens include flavonoids such as estrogenic flavonoids, genistein, daidzein, glycitin, biochanin A, formononetin and equol and mixtures thereof, acetyl



and malonyl esters of genistein and daidzein, and glucosides of genistein and daidzein. It should be noted that the aforementioned list is not exclusive, and may include other phytoestrogens known to persons of ordinary skill in the art.

5                   **DUAL COMPARTMENT PACKAGE**

10                   Compositions which include retinoids are generally unstable and may undergo chemical degradation. Moreover, it has been surprisingly found that boosters, although beneficial for enhancing the retinoid benefits, also contribute to the chemical instability of retinoids. The booster induced retinol destabilization dramatically reduces the overall efficacy of the boosted retinoid composition when both ingredients are contained in a single formula. Therefore, in order to protect against retinoid breakdown while still providing the beneficial effects of retinoid boosters, the present invention provides a dual compartment package that contains a first composition containing retinoids in a first compartment and a second composition containing at least one retinoid booster in a second compartment. The first composition provides a first benefit to the skin while the second composition works to boost or enhance the effect of the first benefit.

20                   As a further retinoid enhancing benefit, phytoestrogens are an essential component of the present invention. Phytoestrogens such as genistein and daidzein synergistically interact with retinoids to deliver skin benefits. However, phytoestrogens contribute to the oxidation, and thus the degradation of retinoids. Therefore, the present invention provides the phytoestrogen as part of the second composition in the second compartment of the dual compartment package, to further enhance the effect of the first benefit.

**J6673(C)**  
**Y2-R566-EDG**

The dual compartment package may be designed in various ways known to persons of ordinary skill in the art as long as the purpose of providing the first and second compositions in two separate containers is achieved. In one embodiment, the dual compartment package is in the form of two jars or bottles adjointly  
5 attached. In a second embodiment, the dual compartment package is in the form of a single bottle/jar with a division separating an interior of the bottle/jar into a first and second compartment. Other embodiments are contemplated as being within the scope of the present invention as long as the compositions are retained separately.

**Cosmetically Acceptable Vehicle**

The product according to the present invention also comprises a cosmetically acceptable vehicle to act as a dilutant, dispersant, or carrier for the  
15 active components in the either or both the first and second compositions, so as to facilitate their distribution when the composition is applied to the skin.

Vehicles other than or in addition to water can include liquid or solid emollients, solvents, humectants, thickeners and powders. An especially preferred  
20 nonaqueous carrier is a polydimethyl siloxane and/or a polydimethyl phenyl siloxane. Silicones of this invention may be those with viscosities ranging anywhere from about 10 to 10,000,000 centistokes at 25 °C. Especially desirable are mixtures of low and high viscosity silicones. These silicones are available from the General Electric Company under trademarks Vicasil, SE and SF and from the Dow Corning  
25 Company under the 200 and 550 Series. Amounts of silicone which can be utilized in the compositions of this invention range anywhere from 5 to 95%, preferably from 25 to 90% by weight of the composition.

### Optional Skin Benefit Materials and Cosmetic Adjuncts

5 In either one or both of the first and second compositions of the present invention, an oil or oily material may be present, together with an emulsifier to provide either a water-in-oil emulsion or an oil-in-water emulsion, depending largely on the average hydrophilic-lipophilic balance (HLB) of the emulsifier employed.

10 Various types of active ingredients may be present in either one or both of the first and second cosmetic compositions of the present invention and are described below. Actives are defined as skin or hair benefit agents other than emollients and other than ingredients that merely improve the physical characteristics of the composition. Although not limited to this category, general  
15 examples include sunscreens, skin lightening agents, tanning agents.

Sunscreens include those materials commonly employed to block ultraviolet light. Illustrative compounds are the derivatives of PABA, cinnamate and salicylate. For example, octyl methoxycinnamate and 2-hydroxy-4-methoxy  
20 benzophenone (also known as oxybenzone) can be used. Octyl methoxycinnamate and 2-hydroxy-4-methoxy benzophenone are commercially available under the trademarks, Parsol MCX and Benzophenone-3, respectively.

25 The exact amount of sunscreen employed in the emulsions can vary depending upon the degree of protection desired from the sun's UV radiation.

**J6673(C)**  
**Y2-R566-EDG**

Another preferred optional ingredient is selected from essential fatty acids (EFAs), i.e., those fatty acids which are essential for the plasma membrane formation of all cells. In keratinocytes EFA deficiency makes cells hyperproliferative. Supplementation of EFA corrects this. EFAs also enhance lipid biosynthesis of epidermis and provide lipids for the barrier formation of the epidermis. The essential fatty acids are preferably chosen from linoleic acid,  $\gamma$ -linolenic acid, homo- $\gamma$ -linolenic acid, columbinic acid, eicosa-(n-6,9,13)-trienoic acid, arachidonic acid,  $\gamma$ -linolenic acid, timnodonic acid, hexaenoic acid and mixtures thereof.

Emollients are often incorporated into cosmetic compositions of the present invention. Levels of such emollients may range from about 0.5% to about 50%, preferably between about 5% and 30% by weight of the total composition. Emollients may be classified under such general chemical categories as esters, fatty acids and alcohols, polyols and hydrocarbons.

Esters may be mono- or di-esters. Acceptable examples of fatty di-esters include dibutyl adipate, diethyl sebacate, diisopropyl dimerate, and dioctyl succinate. Acceptable branched chain fatty esters include 2-ethyl-hexyl myristate, isopropyl stearate and isostearyl palmitate. Acceptable tribasic acid esters include triisopropyl trilinoleate and trilauryl citrate. Acceptable straight chain fatty esters include lauryl palmitate, myristyl lactate, oleyl eurate and stearyl oleate. Preferred esters include coco-caprylate/caprate (a blend of coco-caprylate and coco-caprate), propylene glycol myristyl ether acetate, diisopropyl adipate and cetyl octanoate.

Suitable fatty alcohols and acids include those compounds having from 10 to 20 carbon atoms. Especially preferred are such compounds such as cetyl, myristyl, palmitic and stearyl alcohols and acids.

Among the polyols which may serve as emollients are linear and branched chain alkyl polyhydroxyl compounds. For example, propylene glycol, sorbitol and glycerin are preferred. Also useful may be polymeric polyols such as polypropylene glycol and polyethylene glycol. Butylene and propylene glycol are also especially preferred as penetration enhancers.

Exemplary hydrocarbons which may serve as emollients are those having hydrocarbon chains anywhere from 12 to 30 carbon atoms. Specific examples include mineral oil, petroleum jelly, squalene and isoparaffins.

Another category of functional ingredients within the cosmetic compositions of the present invention are thickeners. A thickener will usually be present in amounts anywhere from 0.1 to 20% by weight, preferably from about 0.5% to 10% by weight of the composition. Exemplary thickeners are cross-linked polyacrylate materials available under the trademark Carbopol from the B.F. Goodrich Company. Gums may be employed such as xanthan, carrageenan, gelatin, karaya, pectin and locust beans gum. Under certain circumstances the thickening function may be accomplished by a material also serving as a silicone or emollient. For instance, silicone gums in excess of 10 centistokes and esters such as glycerol stearate have dual functionality.

Powders may be incorporated into one or both of the first and second cosmetic compositions of the cosmetic product of the present invention. These powders include chalk, talc, Fullers earth, kaolin, starch, smectite clays, chemically modified magnesium aluminum silicate, organically modified montmorillonite clay,

**J6673(C)**  
**Y2-R566-EDG**

hydrated aluminum silicate, fumed silica, aluminum starch octenyl succinate and mixtures thereof.

5 Other adjunct minor components may also be incorporated into one or both of the first and second compositions of the cosmetic product of the present invention. These ingredients may include coloring agents, opacifiers and perfumes. Amounts of these materials may range anywhere from 0.001% up to 20% by weight of the composition.

10 The first and second compositions of the cosmetic product of the present invention are intended primarily as a product for topical application to human skin, especially as an agent for conditioning and smoothening the skin, and preventing or reducing the appearance of wrinkled or aged skin.

15 In use, a small quantity of the first composition, for example from 1 to 5ml, is applied to exposed areas of the skin, from a suitable container or applicator and, if necessary, it is then spread over and/or rubbed into the skin using the hand or fingers or a suitable device. Simultaneously, a small quantity of the second composition, for example from 1 to 5 ml, is applied to exposed areas of the skin, 20 from a suitable container or applicator and, if necessary, it is also spread over and/or rubbed into the skin using the hand or fingers or a suitable device. Therefore, depending upon the intensity of treatment benefits desired, the first and second compositions may be used alone, simultaneously, or in consecutive order.

25

**J6673(C)**  
**Y2-R566-EDG**

**Product Form and Packaging**

The topical skin treatment composition of the invention can be formulated as a lotion, a fluid cream, a cream or a gel.

5

**EXAMPLE 1**

**Methods**

10 Retinol (50% in tween 80) was dissolved in approximately 50% aqueous ethanol to provide a solution giving an OD at 360nm of approximately 0.6 when measured in a 200 µl volume in a 96 well plate using a standard 96 well spectrophotometer.

15 Booster molecules were added at approximately 0.1% concentration and the OD 360 measured as above immediately and after 60 hours at room temperature in the dark. A correction was applied to the OD after 60 hours (divide by 0.85) to account for increased concentration of the retinol due to evaporation of solvent from the plate.

**J6673(C)**  
**Y2-R566-EDG**

Results

TABLE 1

BOOSTER	FOLD INCREASE IN RATE OF RETINOL LOSS
CITRAL	3.1
CITRONELLOL	1.5
COCAMIDE DEA	1.9
COUMARIN	1.4
DAMASCONE	3.7
1,3 DIMETHYL 2 IMIDAZOLIDINONE	1.4
GERANIOL	1.3
18b GLYCERHETINIC ACID	1.6
8 OH QUINOLINE	1.5
N LAURY SARCOSINE	2.6
LINALOOL	2.0
LINOLEAMIDE DEA	3.0
LINOLEIC ACID	3.4
ALPHA IONONE	1.3
LINSEED OIL	1.5

5

The Boosters tested caused marked increases in the instability of the retinol.

This will make it necessary to use formulation/packaging options providing considerably better stability to the retinol when boosters are used compared to those needed for retinol alone.

10



EXAMPLE 2

To establish whether synergistic inhibition of transglutaminase expression occurred by combinations of B1 and B5 active compounds with retinol, it is essential to determine the dose response profiles (including IC50 values) of the active compounds when tested individually in the presence of retinol. This data was used to determine an appropriate sub-maximal inhibitory concentration of each active compound, to make it possible to identify synergistic effects of mixtures of the active compounds in the presence of retinol. In order to demonstrate synergy of two compounds, it is essential to select concentrations to test that are at most IC20, in other words a compound concentration that individually boosts the retinol inhibition of transglutaminase expression by 20%. Two such compounds should have an additive inhibition of 40%. Using this strategy to determine concentration leaves a window of 40-100% for further transglutaminase inhibition for detecting synergy of the two compounds under examination. A more challenging concentration criteria would be selecting concentrations of compounds which alone showed no boosted retinol inhibition of transglutaminase. In this study however we chose an even more challenging criteria. We selected concentrations of compounds that were 10 fold and 100 fold lower than the minimally effective transglutaminase inhibiting concentration. Identification of synergistic combinations using such very low concentrations would mean that the most effective synergistic combinations were identified.

The data in the following table represents the concentrations of compound that are 2 logs lower than the minimally inhibitory compound concentration. These were the concentrations used in the B1/B5 combination studies.

**J6673(C)**  
**Y2-R566-EDG**

5

**TABLE 2**

Compound	Concentration
<b>B1 Compounds</b>	
Linoleoyl monoethanolamide	1.00E-06
Palmitamide monoethanolamide	1.00E-06
Farnesol	3.16E-06
Hexyl sphingosine	1.00E-06
Utrecht-2	3.16E-08
Oleoyl betaine	3.16E-07
Oleoyl hydroxyethylimidazoline	1.00E-08
Cocoyl hydroxyethylimidazoline	1.00E-09
Ursolic acid	1.00E-08
Alpha-ionone	3.16E-05
<b>B5 Compounds</b>	
Ketoconazole	1.00E-09
Miconazole	3.16E-09
Climbazole	1.00E-08
Amino benzotriazole	1.00E-06
3,4-dihydroquinoline	1.00E-06
2-hydroxyquinoline	3.16E-06

To investigate synergistic inhibition of transglutaminase expression by combinations of B1 and B5 active compounds with retinol, selected combinations of compounds were tested at concentrations given in the above table. The following data was obtained:

15

TABLE 3

Combination	B1 Compound	B5 Compound	Mean % control TGase
<b>B1 / B5</b>	Farnesol	Ketoconazole	84%
<b>B1 / B5</b>	Hexanoyl sphingosine	Miconazole	68%
<b>B1 / B5</b>	Hexanoyl sphingosine	Ketoconazole	64%
<b>B1 / B5</b>	Hexanoyl sphingosine	3,4-dihydroquinoline	89%
<b>B1 / B5</b>	Hexanoyl sphingosine	Aminobenzotriazole	81%
<b>B1 / B5</b>	Hexanoyl sphingosine	Climbazole	63%
<b>B1 / B5</b>	Oleoyl betaine	Ketoconazole	81%
<b>B1 / B5</b>	Oleoyl hydroxyethylimidazoline	Climbazole	52%
<b>B1 / B5</b>	Cocoyl hydroxyethylimidazoline	Climbazole	71%
<b>B1 / B5</b>	Ursolic acid	2-hydroxyquinoline	74%
<b>B1 / B5</b>	Alpha-ionone	Miconazole	84%
<b>B1 / B5</b>	Alpha-ionone	Ketoconazole	82%
<b>B1 / B5</b>	Alpha-ionone	2-hydroxyquinoline	76% ✓
<b>B1 / B5</b>	Utrecht-2	Aminobenzotriazole	82%
B1 / B5	Linoleoyl monoethanolamide	Ketoconazole	93%
B1 / B5	Linoleoyl monoethanolamide	Climbazole	94%
B1 / B5	Naringenin	Ketoconazole	100%
B1 / B5	Quercetin	Climbazole	92%
B1 / B5	Castor Oil monoethanolamide	Climbazole	98%
B1 / B5	Castor Oil monoethanolamide	Clotrimazole	100%

5

The efficacy of the B1/B5 combinations splits into two classes – particularly effective combinations (bolded in the above table, i.e., the first fourteen combinations) and barely effective combinations (not bolded, i.e., the last six combinations). It was unexpected that certain B1/B5 combinations performed better than other combinations. Those combinations which were barely effective were (i) fatty acid amides + azoles (ii) hydroxy fatty acid amides + azoles and (iii) naringenin/quercetin + azoles. The effective combinations contained B1 boosters

**J6673(C)**  
**Y2-R566-EDG**

combined with B5 boosters from the following classes: fatty hydroxyethyl imidazoline surfactants, cyclic aliphatic unsaturated compounds, polycyclic triterpenes, n-substituted fatty acid amides.

5

**EXAMPLE 3**

This example shows the synergy of retinoids and phytoestrogens:

(a) Cell culture method:

10

Human adult fibroblasts obtained from sun-protected inner arm of 25-30 year female volunteer were used in this. Cells were grown in 1:1 DMEM/Hams F12 media containing 10% FBS, maintained at 37°C in a 5% CO<sub>2</sub> atmosphere under normal atmosphere oxygen tension. Third passage adult fibroblasts were grown in DMEM media with 10% FBS in 12-well plates at a seeding density of 2500 cells/ml/well. The cells at 80% confluence were rinsed in serum free and phenol red free (PRF) DMEM media twice. Pre-treatment with phyto-compounds for 4 hours was conducted and then dosed with retinoids and was incubated for 48 hours. After the incubation, the wells were washed twice with 1X PBS and the cell monolayer was harvested in 100 µl cell lysis buffer (contains 1X PBS, 1% Triton X, 0.5% sodium deoxycholate, 0.1% SDS containing protease inhibitor (10mg/ml PMSF in isopropanol, 10µl/ml). The suspension was spun at 14000rpm for 10 minutes, the supernatant collected and an aliquot of this supernatant was used for protein quantification. Protein concentration was determined using Pierce protein kit. The remainder of 100 µl supernatant (cell lysate) was denatured in a mixture of 40 µl sample buffer (NOVEX) and 0.5% Beta mercaptoethanol (BME) and by boiling the sample for 5 minutes. Equal amount of protein was then loaded onto

15

20

25

**J6673(C)**  
**Y2-R566-EDG**

16% Tris-glycine gels for protein analysis by SDS page and Western Immunoblotting for CRABP-2 protein expression.

5

(b) Detection of Cellular Retinoic Acid Binding Protein II (CRABP-II):

Within the cells, retinol and retinoic acid are bound to specific cellular binding proteins, 2 of the major proteins are CRABP-I and II (Roos et al., Pharmacological reviews: 50, 315-333, 1998). These proteins act in regulating the intracellular concentration of retinoids by acting as both storage or shuttle proteins in retinoid metabolism. High or low levels of retinoids cause cell damage, including cell death, therefore regulation of constant levels of retinoids and its binding proteins are very critical for cell survival. The levels of this protein are regulated by the amount of retinoic acid within the cells. Higher cellular levels of retinoids increase the expression of CRABP-II. Therefore, the amount of this protein in the cells, is a measure of the retinoid activity of the cells. Skin cells contain high levels of CRABP-II both in the epidermis and the dermis. CRABP-II response to retinoid administration in fibroblasts in vitro is used as a reproducible measure of retinoid bioactivity that predict human skin responses. (Elder et al., J. Invest. Dermatol., 106: 517-521, 1996). Increase in CRABP-II is also associated with increased epidermal differentiation, and dermal retinoid action. Therefore, in these studies we used CRABP-2 expression of fibroblasts as a measure of retinoid activity leading to increased epidermal differentiation (skin conditioning and dry skin benefit) and dermal collagen and extracellular matrix synthesis (anti-aging, anti-wrinkling benefits).

**J6673(C)**  
**Y2-R566-EDG**

To measure the levels of CRABP-II in the fibroblasts, the equal amount of protein of cell supernatant were loaded onto to nitrocellulose blots in a dot blot apparatus as instructed by the manufacturer, and immunostaining was carried out using monoclonal antibodies to CRABP-II according to standard procedures. The CRABP-II protein band was visualized in the Dot Blots using the chemiluminescence system obtained from Santa Cruz Biotechnology (SantaCruz, CA). The bands in the film were quantitated by densitometric scanning, the data from the triplicate samples were calculated as % of control and expressed in the following tables as % increase over control (with control as 100%) +/-SD triplicates.

**EXAMPLE 4**

This example shows the stability of Retinol in the Presence of Phytoestrogenic Flavonoids.

Retinol was dissolved as a 10% solution in aqueous ethanol (1:1 water:ethanol). This solution was diluted to 0.001% approximately 30  $\mu$ M). This solution gave an OD of about 0.35 absorption unit at 360 nm in a 96 well plate spectrophotometer.

Aqueous ethanolic stock solutions of the genistein, daidzein were prepared as 0.1%, 0.01% or 0.001%. To 200  $\mu$ l of 0.001% retinol solution in a 96 well plate was added 20  $\mu$ l of the flavonoid (i.e. 1-10 dilution) giving a final flavonoid concentration of 0.01, 0.001 and 0.0001%. The plates were mixed and an initial OD reading was taken at 360 nm. The plates were incubated at room temperature in the dark for up to 2 days and subsequent readings were taken at

**J6673(C)**  
**Y2-R566-EDG**

8, 24 and 48 hours. The OD readings at these time points were normalized to the 0 time point reading. The retinol stability was expressed as % of retinol (OD reading) at 0 time. The data is shown in example 5.

5

10

EXAMPLE 5

15

In the 2 tables shown below, synergy between genistein and daidzein and retinoids were tested. In both the studies genistein was delivered to the cells in a soluble form in DMSO: ethanol. 1  $\mu$ M genistein alone stimulated CRABP-II significantly. Both genistein and daidzein stimulate retinoid activity in a synergistic manner. All the retinoids tested, except retinyl acetate showed synergy with genistein and daidzein. These data support the our claim that the phytoestrogenic flavonoids genistein and daidzein, when supplied to cells in a soluble form, synergistically enhanced the activity of retinoids.

20

**J6673(C)**  
**Y2-R566-EDG**

**TABLE 4**

Synergy between genistein and retinoids.

	CRABP-II production	% as Control	P value vs. Control p<0.05	P value vs. retinoids p<0.05	Synergy
Control	0.29+/-0.07	100+/-27	1		
10nM Retinoic acid	1.24+/-0.29	428+/-101	0.0055	1	
1nM Retinoic acid	0.97+/-0.47	335+/-162	0.068	1	
100nM Retinyl Linoleate	0.52+/-0.3	181+/-110	0.28	1	
100nM Retinyl Palmitate	1.26+/-0.51	434+/-177	0.032	1	
100NM Retinyl Acetate	0.60+/-0.32	209+/-118	0.19	1	
1µM Genistein	1.9+/-0.71	655+/-247	0.018		
1µM Genistein + 10nM Retinoic acid	4.18+/-0.31	1441+/-108	3.23E-05	2.96E-04	YES
1µM Genistein + 1nM Retinoic acid	4.01+/-0.61	1383+/-394	0.00049	0.012	YES
1µM Genistein + 100nM Retinyl linoleate	4.08+/-1.14	1408+/-213	0.0045	0.000982	YES
1µM Genistein + 100nM Retinyl palmitate	4.32+/-0.13	1489+/-47	160E-06	5.76E-04	YES
1µM Genistein + 100nM Retinyl acetate	2.32+/-0.91	800+/-313	1.80E-02	3.80E-02	NO



**J6673(C)**  
**Y2-R566-EDG**

**TABLE 5**

**Synergy Between Daidzein and Retinoids**

	CRABP-2 production	% as Control	P value vs. Control p<0.05	P value vs. retinoids p<0.05	Synergy
Control	0.29+/-0.07	100+/-27	1		
10nM Retinoic acid	1.24+/-0.29	428+/-101	0.0055	1	
1nM Retinoic acid	0.97+/-0.47	335+/-162	0.068	1	
100nM Retinyl Linoleate	0.52+/-0.3	181+/-110	0.28	1	
100nM Retinyl Palmitate	1.26+/-0.51	434+/-177	0.032	1	
100NM Retinyl Acetate	0.60+/-0.32	209+/-118	0.19	1	
1µM Daizedein	1.49+/-0.66	513+/-227	0.035		
1µM Daizedein + 10nM Retinoic acid	3.42+/-1.01	1181+/-350	0.0059	0.023	YES
1µM Daizedein + 1nM Retinoic acid	3.52+/-0.47	1213+/-163	0.000309	0.027	YES
1µM Daizedein + 100nM Retinyl linoleate	3.29+/-0.14	1136+/-142	0.00024	0.00078	YES
1µM Daizedein + 100nM Retinyl palmitate	2.51+/-0.19	865+/-65	4.90E-05	1.69E-02	YES
1uM Daizedein + 100nM Retinyl acetate	2.27+/-1.4	782+/-489	0.07	0.11	NO

5

**EXAMPLE 6**

Example 5: The following tables show the effect of genistein and daidzein on destabilizing retinol. The experiment was done as described in methods section. The OD readings from duplicate measurements were averaged and given here.

10

J6673(C)  
Y2-R566-EDG

TABLE 6

Retinol stability in the presence of Genistein:

Time(hours)	0% Genistein	0.0001%	0.01%	0.01%
8	92	84	82	82
24	86	76	74	80
48	81	77	73	76

5

TABLE 7

Retinol stability in the presence of Daidzein

Time (hours)	0% Daidzein	0.0001%	0.001%	0.01%
8	92	82	80	73
24	86	78	79	69
48	81	76	76	68

10 Retinol alone in the absence of any agents degraded slowly (8% by 8 hours, 14% by 24 hours and 19% by 48 hours). However, in the presence of genistein and daidzein the degradation of retinol was accelerated. As early as 8 hours, 16-18% of retinol was degraded in the presence of these flavonoids. This suggests that both genistein and daidzein caused marked increases in the instability of retinol. This will make it necessary to use special packaging, one  
15 compartment for retinol and another for the flavonoids in products containing retinoids and the flavonoids.

While the present invention has been described herein with some specificity, and with reference to certain preferred embodiments thereof, those of ordinary skill

**J6673(C)**  
**Y2-R566-EDG**

in the art will recognize numerous variations, modifications and substitutions of that which has been described which can be made, and which are within the scope and spirit of the invention. It is intended that all of these modifications and variations be within the scope of the present invention as described and claimed herein, and that  
5 the inventions be limited only by the scope of the claims which follow, and that such claims be interpreted as broadly as is reasonable. Throughout this application, various publications have been cited. The entireties of each of these publications are hereby incorporated by reference herein.

10

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
221